



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ARITHMETIC.

Conducted by B. F. FINKEL, Kidder, Missouri. All Contributions to this department should be sent to him.

SOLUTIONS TO PROBLEMS.

22. Proposed by E. S. LOOMIS, A. M., Ph. D., Professor of Mathematics, Baldwin University, Berea, Ohio.

A borrows \$1,000 from *B* for 10 years, on which he pays 4% semi-annually.

A immediately loans the \$1,000 to *C* for 10 years, who agrees to pay to *A* \$12½ on the first of each month for 120 mos. or 10 yrs., at which time the whole debt is considered canceled, *C* no longer being, in any way, indebted to *A*. Upon the receipt of each of the \$12½ payments made by *C*, *A* immediately reloans it to *D*, *E*, *F*, etc., upon the same conditions as he loaned the \$1,000 to *C*; at the end of 120 mos. all who are indebted to *A* pay up in full all due him, and he (*A*) pays *B* the principal, all interest having been paid when due.

Query: How many dollars has he in hand?

NO SOLUTION RECEIVED.

23. Proposed by H. C. WHITAKER, Professor of Mathematics, Manual Training School, Philadelphia, Pennsylvania.

A rectangular hall 80 feet long, 40 feet wide and 12 feet high, has a spider in one corner of the ceiling. How long will it take the spider to crawl to the opposite corner on the floor, if he crawls a foot in a second on the wall and two feet in a second on the floor?


I. Solution by Professor J. A. TIMMONS, St. Mary's, Kentucky, and the PROPOSER

It seems to me this problem does not belong to arithmetic. Were the *shortest route* required, arithmetic would solve it; a line from *E* (the *side* wall supposed to be lying down) to *C* being the required distance=95.41 ft. The

time to travel *this* route = $22.02 + \frac{73.4}{2} = 58.72$ seconds.

Were the spider to descend the *end* wall, a line from *G* to *C* would give the shortest distance by that route; but although this distance is *longer* than the other one, being 100.32 ft., the *time* would be shorter, being only $13.085 + \frac{87.233}{2} = 56.691$ seconds. Hence we see that arithmetic alone will not solve it.

Let *F* be point when spider reaches the floor; call *AF* *x*; then *FD* = 80 - *x*.



We have $EF = \sqrt{x^2 + 144}$, and $CF = \sqrt{8000 - 160x + x^2}$; hence the time required in seconds = $\sqrt{x^2 + 144} + \sqrt{2000 - 40x + \frac{x^2}{4}}$; that is, $u = \sqrt{x^2 + 144}$